



Promoting Child Development

Theme 2

INTRODUCTION

Any health supervision encounter with children involves promoting healthy child development. Understanding child development and the application of its principles sets the care of children apart from that of adults. Infants must grow to be children, then adolescents, and then adults. Health promotion to ensure physical, cognitive, and social emotional health as well as to protect the child from infectious diseases and injuries (intentional and unintentional) supports the healthy development of the child. Successful health promotion efforts should take into account the developmental reality of the child now, as well as her developmental expectations for the next months and her developmental potential for growth over time.

Encouraging development of the growing child recognizes the wonder of brain development with its concurrent increases in volume, size, and synapse formation. Physical growth to support brain development is essential. Even more important are the influences of stimulation, social interactivity, family, culture, and community.

The development of the infant, child, or youth with special health care needs is addressed in separate sections within this theme. Even a child whose brain growth and function have been impaired by injury or early neglect has a

developmental potential that must be discerned and supported to achieve the best possible outcome for that child.

Monitoring Child and Adolescent Development

Developmental surveillance and screening of children and adolescents are integral components of health care supervision. Surveillance of children and adolescents is a continuous and cumulative process that is used to ensure optimal health outcomes. For example, it is essential in identifying and treating children with developmental and behavioral problems.

Early identification of children with developmental delay is critical for diagnosing and providing early therapeutic interventions.¹ The parents' report of current skills can accurately identify developmental delay, even though they may not



Long-term outcomes for all infants are improved when health care professionals emphasize the abilities of the infant and facilitate opportunities for the parents to have early physical contact through breastfeeding, rooming-in, holding skin-to-skin, and cuddling the infant.

recognize it as such. Standardized developmental parent-completed questionnaires make it easier for health care professionals to systematically elicit information that is reliable and valid.¹ During all encounters, health care professionals also must listen carefully to parental concerns and observations about a child's development.²

Comprehensive child development surveillance includes:

- Eliciting and attending to the parents' concerns
- Maintaining a developmental history
- Making accurate and informed observations of the child
- Identifying the presence of risk and protective factors
- Periodically using screening tests
- Documenting the process and findings

In monitoring development during infancy and early childhood, ongoing surveillance is supplemented and strengthened by standardized developmental screening tests that are used at certain visits (9 months, 18 months, and 2½ years) and at other times at which concerns are identified.¹ Currently, no comprehensive developmental screening tests exist for use during the middle childhood or adolescent visits. However, several tools have been developed that are useful in screening for particular problems. For example, the Pediatric Symptom Checklist is a psychosocial screen that can be used to identify cognitive, emotional, and behavioral problems.³ The CRAFFT is a 6-item tool that can be used to screen specifically for drug and alcohol use.⁴

Promoting Child Development: Infancy—Birth to 11 Months

The first year of life is a period of neural plasticity and rapid adjustment to stimuli that allow the infant's brain to develop to its maximum potential or not, depending on his experiences.⁵

Long-term outcomes for all infants are improved when health care professionals emphasize the abilities of the infant and facilitate opportunities for the parents to have early physical contact through breastfeeding, rooming-in, holding skin-to-skin, and cuddling the infant.^{6,7}

Developmentally focused anticipatory guidance should include information on growth and development, talking and reading aloud, safety related to the child's developmental abilities and physical capabilities, sudden infant death syndrome (SIDS), coping with the stressors that make infants vulnerable to abuse (eg, infant crying and maternal postpartum depression), and parenting an infant with special or developmental health care needs. Cultural considerations influence parental perspectives about infant temperament and the parental or caregiver role in supporting the infant's self-regulation. The health care professional must try to understand the complex interrelationship of the family's beliefs, values, and behaviors, which affect how a family protects, teaches, and socializes an infant. Parents' perspectives about the needs of their children and whether they view the infant's behaviors as normal or typical for the child's age are equally important considerations. Because families vary in their responses and behaviors, the health care professional must learn about these customs and seek to understand parents' responses and behaviors, even if they differ from those expected in the community context.

Infants With Special Health Care Needs

Most infants are born healthy, but some are born early or at a low birth weight, or have special health care needs. Parents and other caregivers of an infant with special health care needs will need support and guidance in nurturing the infant and fostering family cohesion. Anticipatory guidance should be

structured around the parents' goals and expectations. Specific guidance can include information on growth and development, feeding concerns, specialized health and developmental care needs for the infant, expectations for achieving developmental milestones, and any specific vulnerability that the family will need to know. The health care professional should explore with families their understanding of their infant's health condition, its impact on the family, their expectations on issues such as family supports and care coordination, and their hopes for the child. Additionally, many families may need assistance with referrals, financial assistance, and other types of supports.

The health care professional plays an important role in identifying conditions that place the infant at risk of disability and warrant immediate referral to early intervention services (Box 1). Health care professionals should note those children who require close developmental surveillance and periodic standardized developmental screening to permit the earliest identification of their need for intervention services due to other risk factors. The health care professional also plays an important and continuing role in providing informed clinical opinion in determining the child's eligibility and the scope of services that are needed by the child and family. Care coordination of screening services and follow-up in the context of the medical home are important. Professionals should, however, be aware that some families may not view early intervention as positive (eg, they may see efforts to screen and evaluate as efforts to stigmatize their child, or they may belong to a culture or religion in which differences are tolerated and accepted and are not "fixed").

Developmental surveillance, screening, and observations are important in all aspects of the child's growth and development. Formal developmental evaluation is indicated if any signs of developmental delay exist, if the parents express concern or questions about their

child's development, or if the child is at risk of developmental challenges because of factors such as prematurity or prenatal exposure to alcohol, drugs, or other toxins. Many parents are aware of developmental delays or irregularities before they are told about them by a health care professional. Their concerns must be promptly responded to and appropriate evaluation must be initiated. This evaluation might begin in the primary care office or might result in an immediate referral to an early intervention program or developmental specialist.

Domains of Development

During a child's life, the most dramatic growth—physical, motor, cognitive, communicative, and social-emotional—occurs during infancy. By 1 year of age, the infant has

BOX 1

Program for Infants and Toddlers with Disabilities (Part C of Individuals with Disabilities Education Act)

Children from birth to age 3 years who exhibit delays in development or are at risk are eligible under federal law for early intervention services that will foster age-appropriate development. The Program for Infants and Toddlers with Disabilities (Part C of the Individuals with Disabilities Education Act [IDEA]) assists states in operating a comprehensive, statewide program of early intervention services for infants and toddlers with disabilities, from birth through age 3 years, and their families.⁸ Eligibility criteria can be found at <http://www.nectac.org/topics/earlyid/partcelig.asp>. A diagnosis is not necessary for enrollment in early intervention programs. Children can be on waiting lists for an evaluation while receiving services. Children from the age of 3 years to school age also are eligible for early intervention through the educational system or through developmental services.

The health care professional plays an important role in identifying conditions that place the infant at risk of disability and warrant immediate referral to early intervention services.

When parents provide consistent and predictable daily routines, the infant learns to anticipate and trust his environment.

nearly tripled his birth weight, added almost 50% to his length, and achieved most of his brain weight. By 8 months of age, brain synapses have increased from 50 trillion to 1,000 trillion, and remain there through early childhood.⁹ During the remainder of childhood and adolescence, the brain is actively engaged in developing and refining the efficiency of its neural networks, especially in the prefrontal cortex, the critical brain region responsible for decision making, judgment, and impulse control. This dynamic process of neuronal maturation continues into early adulthood.¹⁰

Outcomes for infants who are prenatally exposed to toxins (eg, alcohol, lead, and illicit drugs) are largely determined not by the degree of exposure but by the quality of the nurturing environment.⁵ Studies on early brain development confirm the importance of positive early experiences in the formation of brain cell connections. These early experiences, especially parent-child interactions, have a significant impact on a child's emotional development and learning abilities.

GROSS MOTOR SKILLS

From birth to the end of the first year of life, major changes occur in the infant's gross motor skills. As tone, strength, and coordination improve sequentially from head to heel, the infant attains head control, rolls, sits, crawls, pulls to stand, cruises, and may even walk by 1 year of age. Delays in gross motor milestones, asymmetry of movement, or muscle hypertonia or hypotonia should be identified and evaluated for early intervention referrals. Within the framework of back-to-sleep guidelines,¹¹ it is important to promote age-appropriate and safe opportunities for tummy-time play to allow young infants to master their early motor skills.

FINE MOTOR SKILLS

Hand-eye coordination and fine motor skills also change dramatically during infancy. These abilities progress from reflexive



grasping to voluntary grasp and release, mid-line play, transferring an object from one hand to the other, shaping the hand to an object, inferior then superior pincer grasp, using the fingers to point, self-feeding, and even marking with a crayon by 1 year of age. Babies should be given opportunities to play with toys and food to advance their fine motor skills.

COGNITIVE, LINGUISTIC, AND COMMUNICATION SKILLS

Environmental factors influence the infant's developing brain significantly during the first year of life. When parents provide consistent and predictable daily routines, the infant learns to anticipate and trust his environment. An infant's brain development is

affected by daily experiences with parents and other caregivers during feeding, play, consoling, and sleep routines.¹²

At birth, newborns already hear as well as adults do, but their responses can be difficult for parents to understand. Newborns should have a screening test for hearing before discharge from the hospital, or should be screened before 1 month of age if not born in a hospital. Thereafter, hearing should be screened regularly and whenever parents express concern about hearing and/or language development. Newborns can recognize their parents' voices at birth. By 3 days of age, they can distinguish their mother's voice.

Newborns also have color vision, can see in 3 dimensions, and can track visually. Close up, they show a preference for the pattern of human faces. Visual acuity progresses rapidly from newborn hyperopia to adult levels of 20/20 vision when the child is 5 to 6 years of age. Newborns copy facial expressions from birth, use the emotional expressions of others to interpret events, and understand and use gestures by 8 months of age. By 8 weeks, babies coo; by 6 to 8 months, they begin to babble with vowel-consonant combinations; and by 1 year, they usually speak a few single words. The normal range for the acquisition of these pre-linguistic skills is broad. Beyond babbling, progress depends in part on the language stimulation a child receives. Children who are talked to and read to frequently by loving parents or caregivers have nearly 300 words at age 2 years, a higher number than often acquired by those who have not had this stimulation.

Reading is important for all children, including infants. Health care professionals should educate parents about how to read to infants, the importance of language stimulation, including singing songs to infants and children, reading to them, and talking to them. They also need to understand the transition from the parent talking about pictures in a book to engaging the child in reciprocally talking and pointing to pictures in a book.

Health care professionals also should identify feeding issues related to oromotor function and coordination because these are integral to early pre-linguistic and later communication skills. Special discussions could be used with parents who are unable to communicate verbally or who have a child with special communication needs (such as a child with a hearing loss) to help the parents support normal language development in their children. Exposure to live language has been shown to have a positive impact on early child development, whereas television screen exposure increasingly shows adverse effects.^{13,14}

Children who live in print-rich environments and who are read to during the first years of life are more likely to learn to read on schedule than children who are not exposed in this way.¹⁵ Giving an age- and culturally appropriate book to the child, along with anticipatory guidance to the parent about reading aloud, at each health supervision visit from 6 months to 5 years, has been shown to improve the home environment and the child's language development, especially in children at socioeconomic risk.¹⁶⁻²¹ Parents should make reading with their children part of the daily routine. Reading together in the evening can become an important part of the bedtime ritual beginning in infancy and continuing for years. Books and reading encourage development in multiple domains and are especially important for cognitive and linguistic development.¹⁷⁻¹⁹ Book-handling skills in young children also reflect fine motor skills, and parent-child reading promotes social and emotional development as well. Reading to a young child is often a source of great warmth and good memories for parents and children alike. Parents can use books in various ways, and health care professionals can emphasize to parents with low or no literacy skills that having conversations with their young children about the pictures in books (ie, interactive

Health care professionals should educate parents about how to read to infants, the importance of language stimulation, including singing songs to infants and children, reading to them, and talking to them.

As parents learn to recognize their infant's behavior cues for engagement and disengagement or distress, and consistently respond appropriately to their infant's needs (eg, being fed when hungry or comforted when crying), babies learn to trust and love their parents.

reading) also is an important way to encourage language development (Box 2).

BOX 2

Promoting Literacy

To help parents promote healthy language and cognitive development in young children, *Bright Futures* recommends anticipatory guidance on reading aloud at every health supervision visit from 6 months to 5 years and encourages giving a book at these visits, whenever possible, especially for children at socioeconomic risk.

Many organizations make books available at low or no cost for distribution. For example, Reach Out and Read (<http://www.reachoutandread.org>) is a national nonprofit organization that promotes early literacy by making books a routine part of pediatric primary care so that children grow up with books and a love of reading.^{18,19} Information on trainings, technical assistance, and start-up funding for these books is available for practices or clinics that are interested in implementing a Reach Out and Read program.

SOCIAL-EMOTIONAL SKILLS

As parents learn to recognize their infant's behavior cues for engagement and disengagement or distress, and consistently respond appropriately to their infant's needs (eg, being fed when hungry or comforted when crying), babies learn to trust and love their parents.

Children with special health care needs may not exhibit the same responses as other children. This difficulty can cause parents to feel inadequate because they cannot discern their child's needs. Helping a family recognize even the small gains their child is making provides support to the family and acknowledges

the progress and growth in their child with special needs.

By 3 months of age, infants may interact differently with different people. At about 8 months, an infant shows social referencing, looking to his parents in ambiguous or unfamiliar situations to figure out how to respond. At about the same age, his capacity to discriminate between familiar and unfamiliar people shows itself as stranger anxiety. By 14 months, he develops enough assurance and communication ability to contain his stranger anxiety and deal successfully with a new person. During the first year, the infant's social awareness advances from a tendency to cry when he hears crying, to attempts to offer food, initiate games, and even take turns by 1 year. As autonomy emerges, babies may begin to bite, pinch, and grab what they want. Health care professionals should tell parents to anticipate these infant behaviors and advise on consistent, appropriate (firm but gentle) responses to redirect the infant's behavior.

Different cultures may have various expectations about the age at which children will achieve socially mediated milestones. It is, therefore, important to ask not only what the child can do but also what the family expects and allows.

Separation Anxiety

Parents need to know that infants as young as 4 to 5 months of age may be anxious, when they are separated from their parents, to meet strangers or even familiar relatives. Even grandparents need to allow the infant to warm up to them before taking the infant from the mother. This anxiety peaks at about 8 months. This is not a rejection but a normal developmental phase.

Providing time for the infant to get to know a new caregiver in the presence of the mother, before separation, is critically important. There must be consistency in this relationship. Transitions will be easier if a child is encouraged to have a special stuffed animal, blanket, or similar favorite object, which she holds on to as an important companion.

Young children use this *transitional object* to comfort them. Transition is often as difficult for the parent as it is for the child. If the parent is going back to work or school and using child care on a consistent basis, the parent often feels a combination of intense longing for the child, intense guilt, and jealousy. The mother is frequently afraid the infant will love the caregiver more than she loves the mother. Parents need to be reassured that they will remain the most important people to their infant's happiness, well-being, and health. The infant may have intense emotions, including crying and irritability, that are saved for times when she is within the safe embrace of her mother. These expressions reflect the intensity of attachment to the mother. Guidance for both the child and parent may be needed to ease transitions and promote healthy adaptations.

Child Care

In recent decades, interest in child care issues has increased as a growing number of women have entered the labor force and as child care arrangements have reflected the changing needs and interests of contemporary American families.²² Regardless of the location or person providing care, young children benefit when they receive high-quality care. Care that fosters children's healthy development should be offered by caregivers who relate consistently to the children; who are available, physically and emotionally, to respond to each child's needs and interests; and who provide care in a clean, safe, nurturing, and stimulating environment. The fewer children cared for by each provider, the better the situation is for the child. Infant care should have no more than 3 children per provider. Parents should ask whether their child care centers adhere to national standards and are accredited by organizations such as the National Association for the Education of Young Children (www.naeyc.org).²³

Developmental Highlights of Infancy

THE INFLUENCE OF CULTURE ON DEVELOPMENT

Health care professionals should understand that what are often considered "milestones" are less "stones" than "markers," and that these markers shift according to upbringing. The timing for acquisition of any developmental task is determined by surveying many infants to determine the range of accomplishment dates. The populations surveyed are typically the population of convenience. So "milestones" must be understood as *normed to a population*. (However, it is important to note that children are still held to the same standards once they reach kindergarten. Therefore, once a child reaches preschool age, developmental differences should be viewed in light of overall population means.) Table 1 presents examples of milestones that are reached at different ages for different ethnic groups.

TABLE 1

Developmental Milestones by Age and Ethnicity, Months²⁴

Task	Anglo	Puerto Rican	Filipino
Eat solid food	8.2	10.	16.7
Use training cup	12.0	17.1	21.9
Use utensils	17.7	26.5	32.4
Eat finger foods	8.9	9.4	9.5
Wean	16.8	18.2	36.2
Sleep by self	13.8	14.6	38.8
Sleep all night	11.4	14.5	32.4
Choose clothes	31.1	44.2	33.1
Dress self	38.2	44.2	39.2
Play alone	25.0	24.8	12.3
Toilet trained (day)	31.6	29.0	20.4
Toilet trained (night)	33.2	31.8	34.2

Helping parents understand their infant's temperament and their own can help them respond effectively to their infant.

SELF-REGULATION

Infants generally are born with unstable physiologic functions. With maturation and sensitive caregiving, physiologic stability, temperature regulation, sustained suck, coordinated suck, swallow, breath sequences, and consistent sleep-wake cycles will improve. During the first year, the infant's ability to self-regulate (eg, transition from awake to sleep) and modulate his behavior in response to stress are influenced by the environment, particularly by the consistency and predictability of the caregivers. The consistency and predictability of infant feedings and encouragement for regular sleep helps establish an infant's diurnal pattern of waking and sleeping. The infant also develops ways to calm himself and expands his ability to selectively focus on a particular activity. Large individual differences exist in self-regulatory abilities. Infants who are born with special health care needs, such as those who are of low birth weight or small for gestational age, or those born to mothers with diabetes or who abused drugs or alcohol during pregnancy, are at particular risk of problems with self-regulation.

A major component of infant health supervision consists of counseling parents about their infant's temperament, colic, temper tantrums, and sleep disturbances. The "goodness of fit" between parents and infant can influence their interaction. Helping parents understand their infant's temperament and their own can help them respond effectively to their infant.

Crying is stressful for families and frustrating for parents. Health care professionals will want to help parents discover calming techniques and understand that a certain amount of crying is inevitable. Parents should consider who they can ask for help if they are having trouble coping or if they fear they might harm their baby.

SLEEP

Parents need guidance on differentiating between active and quiet sleep because they may assume their infant is getting adequate sleep when taken to the mall, taken to a party, or left in a carrier or swing all day. During these times, infants are more apt to be in active sleep. Active sleep alone is not adequate for appropriate rest and often results in a fussy baby. Health care professionals should help parents understand their infant's need for a consistent, predictable, quiet sleep location, including for nap time. Table 2 presents the key characteristics of various infant states. Table 3 lists typical infant sleep patterns.

SUDDEN INFANT DEATH SYNDROME

Sudden infant death syndrome is the sudden, unexplained death of an infant younger than 1 year. Most SIDS-related deaths occur between the ages of 2 and 4 months.²⁵ Numerous studies have identified the following independent risk factors for SIDS:

- Young maternal age
- Maternal smoking during pregnancy (mothers who smoke have an approximately fivefold increase in SIDS risk for their infants)
- Exposure to secondhand cigarette smoke
- Inadequate prenatal care
- Low birth weight or premature birth
- Prone sleep position for infant
- Infant sleeping on a soft surface
- Bed sharing (infant sleeping with parent or other adult)
- An overheated infant
- Male gender

Three of these risk factors are under parental control during infancy: (1) avoidance of cigarette smoke, (2) where the child sleeps and on what type of surface, and (3) in what position he sleeps.

TABLE 2

Key Characteristics of Various Infant States²⁶

Infant States	Characteristics
Quiet sleep	Very difficult to awaken; regular respirations; little movements; may startle
Active sleep	May awaken and go back to sleep; body movements, eyelid movements; irregular respirations
Drowsy	Increasing body movements, eyelid opening; more easily awakened for a feeding but may return to sleep with comforting
Alert	Alert expression, open eyes, surveys surroundings, especially faces; optimum state for feedings
Active alert	Beginning to fuss and show need for a change; if needs are not met, escalates to crying
Crying	Crying that lasts for more than 20 seconds; usually infant can be comforted with holding, feeding, or diaper change; exploring the duration, intensity, and frequency of crying is needed to determine strategies for interventions

TABLE 3

Typical Infant Sleep Patterns and Sleep Location²⁶⁻²⁸

Activities	Birth to 3 Months	3 to 6 Months	6 to 9 Months	9 to 12 Months	12 to 18 Months	18 to 48 Months
Average sleep in 24 h	14 h	13 h	13 h	13 h	12-13 h	12-13 h
Range of sleep in 24 h	12-16 h	12-15 h	10-14 h	10-14 h	12-14 h	12-14 h
Night awakenings	Depends on feeding routine	2-3	1-3	1-2	0-1	0
Number of naps	Depends on feeding routine	2-4 naps/d (am/pm)	2 naps/d (am/pm)	1-2 naps/d (am/pm)	1-2 naps/d	1 nap/d
Length of naps	1-3 h	2-3 h each	1-3 h each	1-3 h each	1-3 h each	1-2 h each
Sleep location	Bassinet or crib in parents' room	Bassinet or crib in parents' room	Crib	Crib	Crib	2-3 y in own bed

Personal experience and beliefs significantly influence a family's acceptance of specific messages regarding infant sleep position and sleep location.

Room Sharing and Bed Sharing

Parent and infant sleeping practices are influenced by custom and family traditions,^{29,30} and these sleep patterns often are among the last traditions to change in immigrant and minority families.

Room sharing, defined as an infant sleeping in the parents' room in a separate sleep space, is a common practice in many cultures worldwide. In many cultures, sharing a room is viewed as a part of the parents' overall commitment to their children's well-being. African American families, for example, view sharing a room as normal, unrelated to perceived infant sleep problems, attachment concerns, breastfeeding, or household crowding.³¹ White parents who share a room with their infants, in contrast, more frequently cite infant sleep problems as the reason for this practice. Increasing evidence shows that room sharing is associated with a reduced risk of SIDS.

Sleep practices in which parents and infants share a bed also are common in many cultures. Bed sharing can take the form of mother, father, and baby together in the same bed, to mother and baby together with father sleeping elsewhere, to all family members in the same bed. Advocates of this practice cite its importance in facilitating breastfeeding, promoting parent-infant attachment, and allowing parents to quickly comfort a fussy infant. Opponents express concerns about safety and that bed sharing promotes an unhealthy attachment between the parent and child and can impede the child's progress toward independence.

A review of the dangers associated with placing children younger than 2 years in adult

beds include overlying by a parent, sibling, or other adult sharing the bed; wedging or entrapment of the child between the mattress and another object; head entrapment in bed railings; and suffocation on water beds or because of clothing or bedding causing oronasal obstruction.^{27,32,33} The issue of parents lying on the infant has received extensive study.³⁰

Sleep Position

Despite the recommendation of supine sleep position for infants, approximately 17% of babies continue to be placed in the prone position for sleep.³⁴ Prone sleep position is used more often by African American families and in child care settings; it is a contributing factor to the disparate SIDS rates. Side lying, an alternative sleep position that is practiced by many families who are concerned about using the prone position, carries a twofold higher risk for SIDS because of the significant probability that the infant will roll from the side position to the prone position during sleep.

Infants who are accustomed to sleeping supine who are placed in the prone position or on their side to sleep are at higher risk for sudden infant death than infants who usually sleep in the prone or side position.³⁵ The risk of death due to SIDS is 7 to 8 times higher for infants who are put to sleep on their side or in the prone position if they typically are placed in a supine position for sleep.³⁵

Reducing Sudden Infant Death Syndrome Risks

Personal experience and beliefs significantly influence a family's acceptance of specific messages regarding infant sleep position and sleep location.^{36,37} The health care professional should learn the family's views about infant sleep, room sharing, and bed sharing to appropriately tailor SIDS prevention and risk reduction counseling. The American Academy of Pediatrics (AAP) Task Force on Infant Sleep Position and Sudden Infant Death Syndrome



reviewed the evidence and compiled the following recommendations to reduce the risk of SIDS¹¹:

- Supine sleep position is safest for every sleep; side sleeping is not advised.
- Use a firm sleep surface.
- Avoid placing soft objects and loose bedding in cribs, bassinets, and playpens.
- Do not smoke during pregnancy.
- Do not allow smoking in the child’s environment.
- A separate but nearby sleep environment is safest for the infant.
- Avoid overheating the infant; do not over bundle the infant or set the room temperature too high.
- Use of home monitors does not prevent SIDS.

- Parents or other caregivers should not share a bed with the infant if they smoke or are using drugs or alcohol or taking medications that cause drowsiness or fatigue or induce a deep sleep.
- Parents and caregivers should not sleep with their infant on a sofa, couch, or water bed.

DISCIPLINE, BEHAVIORAL GUIDANCE, AND TEACHING

The interaction between the parents and their infant is central to the infant’s physical, cognitive, social, and emotional development, as well as her self-regulation abilities. The infant brings her temperament style, physical attentiveness, health, and vigor to this interaction.

Parents need to understand the differences among discipline, teaching, and punishment so that they can introduce appropriate measures for correcting and guiding their infant’s behavior.

DEVELOPMENTAL MILESTONES AT A GLANCE — INFANCY ³⁸				
Age	Gross Motor	Fine Motor	Cognitive, Linguistic, and Communication	Social-Emotional
2 Months	Head up 45° Lift head	Follow past midline Follow to midline	Laugh Vocalize	Smile spontaneously Smile responsively
4 Months	Roll over Sit—head steady	Follow to 180° Grasp rattle	Turn to rattling sound Laugh	Regard own hand
6 Months	Sit—no support Roll over	Look for dropped yarn Reach	Turn to voice Turn to rattling sound	Feed self Work for toy (out of reach)
9 Months	Pull to stand Stand holding on	Take 2 cubes Pass cube (transfer)	Dada/Mama, nonspecific Single syllables	Wave bye-bye Feed self

KEY

Black Color: 50% to 90% of children pass this item.

Green Color: More than 90% of children pass this item.

These norms are taken from the DENVER II, and are based upon the administration and interpretation as set forth in the DENVER II Training Manual (copyright 1992).

These milestones are provided as a reference only. Reference to these milestones does not take the place of a standardized measurement of healthy child development or discourage a developmental discussion with a health care provider.

Parental concerns are highly accurate markers for developmental disability, and it is essential for the health care professional to be sensitive to these concerns.

Parents need to understand the differences among discipline, teaching, and punishment so that they can introduce appropriate measures for correcting and guiding their infant's behavior. It is important to discuss distraction as a developmentally appropriate discipline for infants. It also may be beneficial to discuss strategies to prevent the need for disciplinary measures by avoiding overtiredness through consistent daily routines for feeding and sleep and by providing a developmentally appropriate safe home environment.

Parents' ability to respond appropriately to their child's behavior is determined by their own life stresses, their past experiences with other children, their knowledge, their temperament, their own experiences of being nurtured in childhood, and other responsibilities, such as other children in the household, work, and daily household tasks. Their perceptions of the infant also can influence the interaction. These perceptions come from their own expectations, needs, and desires, as well as from the reaction of other people to the child.

The infant's emotions also can be affected by the emotional health of the caregivers. Depression is common in many mothers of infants and can seriously impair the baby's emotional and even physical well-being. Babies of depressed mothers show delays in growth and development, diminished responsiveness to facial expressions, reduced play and exploratory behaviors, and decreased motor skills.³⁹⁻⁴¹ Parental substance abuse can have similar negative effects. Health supervision for the child must include monitoring the emotional health of the parents or primary caregivers. The health care professional should recognize and provide assistance if parents demonstrate or acknowledge their difficulty in responding to their infant's needs.

Promoting Child Development: Early Childhood—1 to 4 Years

At the beginning of this developmental period, a child's understanding of the world, people, and objects is bound by what he can see, hear, feel, and manipulate physically. By the end of early childhood, the process of thinking moves beyond the "here and now" to incorporate the use of mental symbols and the development of fantasy. For the infant, mobility is a goal to be mastered. For the active young child, it is a mechanism for exploration and increasing independence. The 1-year-old child is beginning to use the art of imitation in her repetition of familiar sounds and physical gestures. The 4-year-old child has mastered most of the complex rules of the languages that are spoken in the home and can communicate thoughts and ideas effectively.

The toddler is beginning to develop a sense of himself as separate from his parents or primary caregivers. By the end of early childhood, the well-adjusted child, having internalized the security of early bonds, pursues new relationships outside of the family as an individual in his own right. Understanding and respecting this evolving independence is a common parental challenge.

Young Children With Special Health Care Needs

Health care professionals who take care of children between the ages of 1 and 4 years have a responsibility to diagnose special health care needs. Because children in this age group grow and progress rapidly, parents anticipate and analyze how their child is reaching developmental milestones such as walking, talking, and socializing. When parents express concerns about how their child is developing, the health care professional should listen and observe carefully. A "wait and see" attitude will not suffice, particularly if the child falls into an at-risk group. A proactive approach is essential.

Several tools are available for identifying a child with special health care needs. Parental concerns are highly accurate markers for developmental disability, and it is essential for the health care professional to be sensitive to these concerns. If developmental delay or disability is suspected, a referral should be made to an appropriate early intervention program or developmental specialist for evaluation. If significant developmental delay or disability is confirmed, the child should be referred to an early intervention program that is matched to the child's and family's needs. With the appropriate services in place, the primary health care professional provides a medical home for the child and, in partnership with the family, assists with ongoing care planning, monitoring, and management across agencies and professionals. The primary care practice team carries out these activities by providing care coordination services. Complicating factors, such as family finances, access to resources, parental health and well-being, and sibling issues, also should be considered. Families whose young children have special health care needs usually find that referrals to parent-to-parent support programs are helpful. (For more information on this topic, see the Promoting Community Relationships and Resources theme.)

Domains of Development

GROSS AND FINE MOTOR SKILLS

The physical abilities of children in the 1- to 4-year age range vary considerably. Some are endowed with natural grace and agility; others demonstrate less fine-tuning in their physical prowess, yet they “get the job done.” As a fearless and tireless explorer and experimenter, the toddler is vulnerable to injury, but appropriate adult supervision and a physically safe environment provide the child with the freedom to take controlled risks.

Many children do not live in safe environments. Parents may try to provide a safe

environment within the confines of their own dwelling, but the immediate community may be characterized by violence, substandard housing conditions, overcrowding, or residence in a shelter. Health care professionals who are aware of these circumstances can better support parents' efforts to find developmentally appropriate surroundings and experiences that allow their children to safely develop their motor skills.

COGNITIVE, LINGUISTIC, AND COMMUNICATION SKILLS

Young children learn through play. If the toddler experienced nurturing and attachment during infancy, she now has a strong base from which to explore the world. The self-centered focus of the young child is related less to a sense of selfishness than to a cognitive inability to see things from the perspective of others. The child's growth in understanding the world around her is evidenced by her linguistic development (ie, by her capacity for naming and remembering the objects that surround her and her ability to communicate her wishes and feelings to important others).

Young children live largely in a world of magic; they often have difficulty differentiating what is real from what is make-believe. Such fantasies, unless scary to the child, are to be expected and encouraged at this stage of development. Some children have imaginary friends. Many children engage in elaborate fantasy play. Learning to identify the boundaries between fantasy and reality and developing an elementary ability to think logically are 2 of the most important developmental tasks of this age.

Parents and other caregivers need to provide a safe environment for these young learners to explore. Children need access to a variety of tools (books and toys) and experiences. They need opportunities to learn through trial and error, as well as through planned effort. Their seemingly endless string

Their seemingly endless string of repetitive questions can test the limits of the most patient parents. These queries, however, must be acknowledged and responded to in a manner that not only provides answers but also validates and reinforces the child's curiosity.

Exposure to books and reading aloud during the time that precedes the formal teaching and learning of reading is central to language development.

of repetitive questions can test the limits of the most patient parents. These queries, however, must be acknowledged and responded to in a manner that not only provides answers but also validates and reinforces the child's curiosity.

The development of language and communication during the early childhood years is of central importance to the child's later growth in social, cognitive, and academic domains. Communication is built upon interaction and relationships. The greater the nurturing and the stronger the connection between parents and child, the greater the child's motivation to communicate will be, first with gestures and then with spoken language. Interactive play and reading are wonderful forums for language enhancement.

Language

Language development usually is described in 3 separate categories: (1) speech (ie, the ability to produce sound, a concept that encompasses rhythm, fluency, and articulation); (2) expressive language (ie, the ability to convey information, feelings, thoughts, and ideas through verbal and other means, including facial expressions, hand gestures, and writing); and (3) receptive language (ie, the ability to understand what one hears and sees). Children can have problems in one area but not in another. Exposure to books and reading aloud during the time that precedes the formal teaching and learning of reading is central to language development. Typical expressive and receptive language acquisitions in the early years include the following:

- **Between the ages of 12 and 18 months**, children make the leap from sound imitation and babbling to the acquisition of a few meaningful words (eg, Dada, Mama, mine, shoe). Through repeated use, these first words teach them how words are used in communication. At the same time that the child gains expressive language, he also shows increased comprehension of simple commands (eg, "say bye-bye") and the names of familiar people and objects. Toddlers expand their communicative repertoire through a variety of gestures (eg, pointing, waving, and playing "pat-a-cake") with and without vocalizations. The child's demonstration of "communicative intent" or proto-declarative pointing (ie, pointing to a desired object and watching to see whether the parent sees it) is an indication of normal social and language development. The absence of pointing and establishing joint attention is a red flag and merits screening for autistic spectrum disorder. At about 18 months of age, most toddlers have begun a word-learning explosion, acquiring an understanding, on average, of 9 new words every day. This pattern continues throughout the preschool years.
- **Between the ages of 18 months and 2 years**, children recognize many nouns and understand simple questions. By the age of 2 years, the expressive language of most children includes 2-word phrases, especially noun-verb combinations that indicate actions desired or observed (eg, "drink juice," "Mommy give").
- **Between the ages of 2 and 3 years**, children usually are speaking in sentences of at least 4 to 5 words. They are able to tell stories and use "what" and "where" questions. They have absorbed the rules for regular plural word forms and for the use of past tense. Their



speech can still be difficult for a non-family member to understand, but it becomes increasingly clear after 3 years of age. A good rule of thumb for normal development is that 75% to 80% of a 3-year-old's speech should be intelligible to a stranger.

- **Between the ages of 3 and 4 years,** children are learning fundamental grammar rules. They have a vocabulary that exceeds 1,000 words, and their pronunciation should be generally understandable. They frequently ask “why” and “how” questions. Their exuberant use of language in play and social interaction often suggests a process of “thinking out loud.”

Parents may ask health care professionals about the effects of being raised in a bilingual home. They can be reassured that this situation permits the child to learn both languages simultaneously as though each language was the mother tongue. If the child is experiencing language delays, however, a consistent language that is spoken by all caregivers may be preferred.

Many aspects of language development seem to be robust in that they develop normally despite environmental conditions. Certain aspects, notably vocabulary and language usage, however, depend heavily on the family and early school experiences if the child is to become proficient.⁵ Thus, the young child who is exposed to an everyday environment that is rich in language through stories, word games, rhymes and songs, questions and conversation in the family and during play, and books will be well prepared for the language-laden world of school. (For more information on this topic, see the Literacy section of this theme.)

Often, hearing loss is first identified as a language delay. If hearing impairment or language delay is a concern, an audiological evaluation is recommended and a referral should be made to early intervention services to optimize language development.

SOCIAL-EMOTIONAL SKILLS

Temperament and Individual Differences

The temperamental differences that were manifested in the feeding, sleeping, and self-regulatory behaviors of the infant are transformed into the varied styles of coping and adaptation by the young child. Some young children appear to think before they act, whereas others are impetuous. Some children are slow to warm up to other people; others are friendly and outgoing. Some children accept limits and rules more easily than others. Some children are highly reactive to changes in their environment and to sensory experiences of all kinds, whereas others are less reactive. Some children tend to express themselves loudly and intensely; others are quieter. Thus, the range of normal behavior is broad.

Understanding the unique temperament profile of the child will better prepare the health care professional to assist parents and other caregivers in understanding the child's behavior, especially when the child's behavioral reactions are confusing or problematic. Discussing with parents how the child's behavior is interpreted within the family, and counseling them when concerns or conflicts emerge between the child's temperament and the caregivers' personal styles, may prevent significant problems later on.

Culture

The culture of the family and community provide a framework within which the socialization process unfolds. Children are heavily influenced by the culture, opinions, and attitudes of their families as they are “taught to act, believe, and feel in ways that are consistent with the values of their communities.”⁴² Culture influences the roles of parents and extended family members in child-rearing practices and the ways in which parents and other adults interact with children. Cultural groups approach parenting in different ways. In some cultures, the mother is expected to be primarily responsible for all aspects of an

Children are heavily influenced by the culture, opinions, and attitudes of their families as they are “taught to act, believe, and feel in ways that are consistent with the values of their communities.”

Ultimately, healthy social and emotional development depends on how children view themselves and the extent to which they feel valued by others.

infant's or toddler's care. In other cultures, the care and nurturing of children is shared among mother, father, and extended family, including aunts, uncles, grandparents, and cousins. This wide circle of caregivers also may have responsibility for disciplining and making other decisions about a child's upbringing.

The increasingly self-aware young child grapples with complex issues, such as gender roles, peer or sibling competition, cooperation, and the difference between right and wrong within this cultural milieu. Aggression, acting out, excessive risk taking, and anti-social behaviors can appear at this time. Caregivers need to respond with a variety of interventions that set constructive limits and help children achieve self-discipline. Fun-filled family activities, such as playing games, reading, vacations, or holiday gatherings, serve as reminders of the joy and laughter the child brings to all. Ultimately, healthy social and emotional development depends on how children view themselves and the extent to which they feel valued by others. The quality of the parent-child relationship is the foundation for emotional well-being and the emerging sense of mastery and self-esteem.

Developmental Highlights of Early Childhood

SELF-REGULATION AND DAILY LIVING TASKS

During the early childhood years, the relative dominance of biologic rhythms is reduced through the development of self-control. Satisfactory self-control allows children to respond appropriately to events in their lives through delaying gratification until important facets of the situation are considered, modulating their responses, remaining calm, focusing on the task, recognizing that their responses have consequences, and behaving in the expected manner to comply with rules and expectations established by their significant caregivers.^{43,44} Usually, these behaviors begin to manifest by 2 to 3 years of age.

Children with inadequate self-control can be impulsive or hyperactive, heightening concerns for safety. At the opposite extreme, children with excessive self-control tend to be anxious or have fixed behaviors. Of course, behavior varies so that a child may exhibit a great variety of behaviors at any given time in response to the same external cues.

Mastering activities in daily life shows that the child is moving toward achieving self-control. Chief among these are learning how to calm herself (which is needed to establish a regular sleep pattern), feed herself, toilet train, and take the major step of attending school. Health care professionals should actively prepare parents and their toddlers for achieving these milestones through discussing these topics and, when concerns persist after counseling, make referrals for appropriate consultation.

SLEEP

By the end of the first year of life, most children should be able to sustain or return to sleep throughout the night, and most parents should allow children to regulate their own nighttime sleep patterns. A consistent bedtime routine that promotes relaxation (eg, bath, book, or song) and the use of a transitional object are extremely helpful. Toddlers and preschoolers generally sleep 8 to 12 hours each night. Exact duration of nighttime sleep varies with the child's temperament, activity levels, health, and growth. The duration and timing of naps will affect nighttime sleeping. Most children awaken from sleep at times during the night, but can return to sleep quickly and peacefully without parental intervention. Sleep problems sometimes reflect separation fears on the part of both parents and children. Parents who feel especially anxious, depressed, or frightened can be reluctant to permit their young child to exercise self-control over sleep patterns at night. Children from 1 to 4 years of age should be allowed to sleep through the night

without a nighttime feeding. Dreams and nightmares can accompany active stages of sleep beginning at these ages. At such times, children may require reassurance that they are protected from the dangers that stir their imagination and intrude upon their calm sleep. Changes, such as acute illness, birth of siblings, and visits from friends and relatives, also can interfere temporarily with established sleep routines. Disorders, such as obstructive sleep apnea, and parasomnias, such as sleep-walking, can begin during these early years, and health care professionals should consider such a possibility in any child who has persistent sleep difficulties.

If health care professionals ask about sleep patterns at each of the visits during early childhood, they will gain rich insights into the child's and family's development. When parents have concerns about their child's sleep, the health care professional should explore, in more depth, the child's daytime behavior, temperament, and mood, as well as events, experiences, conditions, and feelings of family members. Although most issues lend themselves to open dialogue and counseling within the primary care relationship, some conflicts may require further exploration and intervention by a developmental-behavioral or mental health specialist.

TOILET TRAINING

For a child to successfully toilet train, he must have the cognitive capacity to respond to social cues and the neurologic ability to respond to bowel and bladder signals. Parents often want advice about when and how to toilet train a child. The first discussion about toilet training is best introduced at around the 18 Month Visit. Such early counseling can prevent harmful battles between the parents who might be focused on early toilet training and the child who is not yet physically or cognitively ready. In-depth discussion usually begins at the 2 Year Visit. The health care professional should explore the

parents' thoughts about this task and provide guidance to fill in the gaps.

Control of urination and bowel movements is a major step forward in developmental integrity. Successful completion of this task is a source of pride and respect for both the child and the parents.

Daytime control usually is achieved before nighttime dryness. Bed-wetting (nocturnal enuresis) is a common disorder with many possible therapies.⁴⁵ It is much more common in boys and deep sleepers. Enuresis should be investigated if a child continues to wet the bed after age 7 years, if bed-wetting results in problems within the family, or if infection or anatomical abnormalities are suspected. Fortunately, with time, the majority of all children with enuresis develop nighttime urination control.⁴⁶ Bowel control is usually completely achieved by age 3 years.

SOCIALIZATION

When provided the opportunity, toddlers and preschoolers acquire socialization skills and the ability to appropriately interact with other children and adults. Social interaction in early childhood promotes comfort and competence with relationships later in life. The social competencies are developmental assets^{47,48} and, therefore, should be encouraged in children of these ages. Social competencies include planning and decision making with others, positive and appropriate interpersonal interactions, exposure to other cultures and ethnicities, behavioral resistance to inappropriate or dangerous behavior, and peaceful conflict resolution. Young toddlers will observe these behaviors in others, and preschoolers will begin to practice them. They also are inclined to internalize positive or negative attitudes toward themselves and others. Children note differences between groups of people (eg, they express understanding of racial identity as early as 3 years⁴⁹), but they do not ascribe a value; they learn that from the adults in their environments. Opportunities for social interaction can be encouraged in the home with visitors, in play

For a child to successfully toilet train, he must have the cognitive capacity to respond to social cues and the neurologic ability to respond to bowel and bladder signals.

...the health care professional plays a crucially important role in helping parents understand the meaning of their child's behavior, and in assessing the developmental readiness of the child to absorb new lessons about behavioral expectations.

groups, in faith-based organizations, and in public places, such as the park, child care, or preschool.

DISCIPLINE, BEHAVIORAL GUIDANCE, AND TEACHING

Discipline is one tool parents can use to help modify and structure a child's behavior. It encompasses both positive reinforcement of admired behavior (eg, praise for picking up toys) and negative reinforcement of undesirable behavior (eg, a time-out for fighting with a sibling). The eventual incorporation of a functional sense of discipline that reinforces social norms is critical to the child's development. Although often thought of in negative terms, good discipline helps a child fit into the daily family schedule and makes childhood and child rearing pleasant and fun.

Family structure, values, beliefs, and cultural background influence approaches to behavioral guidance and teaching. Health care professionals should discuss with the parents how they were disciplined, how that discipline made them feel, and the most and least effective methods of discipline. In all families and cultures, discipline is a process whereby caregivers and other family members teach the young child, by instruction and example, how to behave and what is expected of her. What the child learns at this stage, and how the parent-child interactions surrounding discipline take form, can have long-term effects on the child's and family's development.

Exploring the roles that siblings play in development also should be addressed. The methods parents use to guide siblings in helping to raise the other family members should be reviewed. The special requirements of children and youth with special health care needs and foster care or adopted children are best discussed openly with all the family members, so that everyone is aware of parental expectations.

Although parents often look to the health care professional as a resource for developing

strategies related to behavioral guidance and teaching, many cultures also look to family, particularly elders, for such guidance and teaching. In most cases, discussions with parents regarding behavioral guidance should explore the parents' goals for the child, as well as the meaning behind the behaviors they wish to modify. Consideration of the child's developmental capacities and temperament profile should be a key component of this discussion. For instance, parents of a 2-year-old child frequently overestimate the child's capacity to integrate rules into everyday behavior, based on their observations of the child's growing understanding of language. With respect to temperament, parents can misinterpret a child's intense and reactive responses as intentionally oppositional rather than as part of her inborn behavioral style. Through explaining these developmental attributes, the health care professional plays a crucially important role in helping parents understand the meaning of their child's behavior, and in assessing the developmental readiness of the child to absorb new lessons about behavioral expectations.

Discussion of discipline is a high priority for the Bright Futures 15 Month and 18 Month Visits because it is important, for later child development, to establish a positive and successful foundation of parent-child interactions regarding behavior. Established negative behaviors can be extremely difficult to change, and, without help, many parents are not able to see the long-term effects of both



their child's behavior and their own choices in guiding them.

At times, the behavior of the child pushes all parents to their emotional limits. Health care professionals should remind parents to avoid meeting the child's anger with their own anger; this reaction teaches the wrong lesson. Many adverse behaviors, such as aggressive acts in the school-aged child have their roots in behavior established in early childhood. Maintaining a sense of humor and taking time away can help parents deal with stressful events. Discussing dilemmas and sharing frustrations with other involved adults are important in maintaining a sense of perspective and humor during difficult periods with the young child.

General features of effective behavioral guidance include several essential components, all of which are necessary for successful discipline⁵⁰:

- A positive, supportive, loving relationship between the parents and child (children want to please their parents)
- Clear expectations communicated to the child in a developmentally appropriate manner
- Positive reinforcement strategies to increase desired behaviors (eg, having fun with the child and other family members sets the stage to reward and reinforce good behaviors with time together in enjoyable activities)
- Removal of reinforcements or use of logical consequences to reduce or eliminate undesired behaviors

Parents can increase the likelihood of achieving their behavioral goals for their child by establishing predictable daily routines and providing consistent responses to their child's behavior. Especially during early childhood, consequences should be administered within close temporal proximity to the target behavior and, if possible, related to the behavior (eg, bring the child in from playtime if she is

throwing sand when asked not to).⁵⁰ Some families (eg, first-time parents or adolescent parents) experience pressure from elders to use harsh or physical means of punishment. Culturally, it may be inappropriate to ignore what an elder has proposed. Parents may feel conflicted when they attempt to use new or different methods of discipline that are not supported within their families or communities.

Parents can use the following techniques to help foster good behavior in their child:

- Praise the child frequently for good behavior. Specific acknowledgement (rather than global praise) helps teach the child appropriate behaviors (eg, "Wow, you did a good job putting that toy away!" rather than "Great!"). Time spent together in an enjoyable activity is a valuable reward for desired behavior.
- Communicate expectations in positive terms. By noting when the child is doing something good, parents will help the child understand what they like and expect. Words such as, "I like it when you play quietly with your brother," or "I like that you climb into your car seat when I ask you to," are nonjudgmental statements and communicate to the child that these are behaviors the parents like.
- Model and role-play the desired behaviors.
- Prepare the child for change in the daily routine by discussing upcoming activities and expected behaviors.
- State behavioral expectations and limits for the child clearly and in a developmentally appropriate manner. These expectations should be few, realistic, and consistently enforced.
- Allow the child time for fun activities, especially as a reward for positive behaviors.
- Remove or avoid the places and objects that contribute to unwanted behavior.

Parents can increase the likelihood of achieving their behavioral goals for their child by establishing predictable daily routines and providing consistent responses to their child's behavior.

Literacy begins in infancy, when parents and other caregivers talk to their baby, and, in early childhood, when toddlers learn to communicate through language, explore their world through imaginary play, and listen to stories, whether read from books or spoken in an oral tradition.

- Use time-out or logical consequences to deal with undesirable behavior.
- Promote consistent discipline practices across caregivers.
- Avoid responding to the child's anger with anger; this reaction teaches the wrong lesson and may escalate the child's response.
- Take time to reflect on their own physical and emotional response to the child's behavior so that they can choose the most appropriate discipline technique.

Conventional disciplinary methods do not work well with children with certain physical or developmental conditions. The following examples illustrate the point that "one size does not fit all" with respect to behavioral guidance:

- Children with poor communication skills often use behavior as a means of communication; caregivers should make every effort to help them develop more effective communication skills.
- Children who have hyperacute responses to their sensory environment require proactive interventions.

Because corporal punishment is no more effective than other approaches for managing undesired behavior in children, the AAP recommends that parents be encouraged and assisted in developing methods other than spanking in response to undesired behavior (Box 4).⁵⁰ Other forms of corporal punishment, such as shaking or striking a child with an object, should never be used. Referral to high-quality parenting programs and counseling should be considered for children with difficult behavioral problems.

LITERACY

Learning to read and write is a complex process that takes time. It requires that children have good, consistent relationships with caring adults who provide one-on-one

interactions and who support the development of oral language. Literacy begins in infancy, when parents and other caregivers talk to their baby, and, in early childhood, when toddlers learn to communicate through language, explore their world through imaginary play, and listen to stories, whether read from books or spoken in an oral tradition. Because young children are active learners, they find joy in exploring and learning the meaning of language and communicating in increasingly sophisticated ways as they move toward literacy.

Parents' and health care professionals' expectations for a young child's literacy accomplishments should be based on developmentally appropriate activities, such as the encouragement of talking, singing, and imaginative play; simple art projects; easy access to books; and frequent reading times. Reading and writing are so linked to development, relationships, and environment that children will vary greatly in when and how

BOX 4

Discipline: Key Messages for Parents

- Discipline means teaching, not punishing.
- All children need guidance, and most children need occasional discipline.
- Discipline is effective when it is consistent; it is ineffective when it is not consistent.
- Parents' discipline should be geared to the child's developmental level.
- Discipline is most effective when the parent can understand the child's point of view.
- Discipline should help a child learn from his mistakes. The child should understand why he is being disciplined.
- Disciplinary methods should not cause a child to feel afraid of his parents.
- A parent should not physically discipline a child if the parent feels out of control.

they learn to read and write. This is true for other complex skills as well.

The National Research Council⁵¹ has identified phonological awareness, shared book reading, and speech-to-print connection as key concepts underlying early literacy. Health care professionals can support literacy by encouraging parents to tell stories, create or visit environments filled with books, find a place at home for imaginary play and art projects, ask their child questions and invite him to talk about his ideas, model literacy behavior by reading newspapers or books daily, and set aside quiet times each day for reading with their child (eg, just before bed). By encouraging parents at every health supervision visit to find age-appropriate ways of incorporating books and reading aloud into children's daily routines, the health care professional can give parents a way to help their children grow up associating books with positive parental attention. These discussions also can help parents understand the role that child care and preschool programs play in helping children get ready to read and write.⁵²

The health care professional's office should reflect reading as a priority, with a specific area set aside to encourage imaginative play, a place with a collection of quality books and magazines where children can look at books or be read to, and a place with information about community libraries and adult and family literacy opportunities. (For more information on this topic, see Box 2 of this theme.) The Reach Out and Read Program has increased the likelihood that parents will read to their children even among families at risk due to low-literacy among parents.^{20,21,53} By giving a book at every health supervision visit from 6 months to 5 years, especially to children at socioeconomic risk, the health care professional can increase the frequency of parental reading aloud, improve the home environment, and help parents increase children's language development.^{20,21}

PLAY

A hallmark of the passage through early childhood is the emergence and steady elaboration of play activities. For the young toddler, play centers on direct explorations into the surrounding world, including the manipulation of objects to create interesting outcomes (eg, the sounds that banging a pot may produce or the interesting results of pouring water into a sandbox). With the development of language, from around age 18 months, play becomes progressively more reflective of the child's remembered experiences and imagined possibilities, as enacted through symbolic play. Thus, a doll comes to represent a living, imaginary person who can be fed, bathed, or scolded—just as the young child has personally experienced in real life.

In *representational* or *symbolic play*, which usually is evident by 2 years of age, the child has a new way of "replaying" the events in her life. Unlike real life, play allows her to control the events and their outcomes. Challenging experiences can be better understood through their re-creation as play. Play can enable the child to better cope with stressful experiences by "taking charge" and developing a preferred story. Confusing or difficult experiences can be mastered through practice in experimentation and planning that play permits.

Many children at this age become attached to *transitional objects* and use them to help them fall asleep, comfort them when they are hurt or upset, and join them in their world of make-believe. The transitional object is a prime example of how the child's active imagination plays a central role in development toward independence and self-regulation.

From 3 to 5 years of age, the child's developmental gains in language and speech, cognitive ability, and fine and gross motor skills allow for increasingly complex forms of play. Play becomes an important modality for practicing and enhancing a broad range of skills,

With the development of language, from around age 18 months, play becomes progressively more reflective of the child's remembered experiences and imagined possibilities, as enacted through symbolic play.

Play provides a window into many aspects of the child's developmental progress and into how she is attempting to understand the events, transitions, and stresses of everyday life.

such as the motor skills and spatial understanding that comes with building with blocks or working with puzzles.

Play is a critical part of development, and toys are a critical part of play. Health care professionals often are asked to recommend appropriate toys for their patients. Toys should be educational and should promote creativity. Parents and health care professionals should avoid toys that make loud or shrill noises, toys with small parts, loose strings, cords, rope, or sharp edges, and toys that contain potentially toxic materials. Toys that promote violence, social distinctions, gender stereotypes, or racial bias also should be avoided. Video games are not recommended for young children, but, if used, they should be screened for inappropriate content. Health care professionals should advise parents on distinguishing between safe and unsafe toys, choosing toys that help promote learning, and using books and magazines to read and play together.⁵⁴

Play provides a window into many aspects of the child's developmental progress and into how she is attempting to understand the events, transitions, and stresses of everyday life. Parents and other caregivers should recognize the importance of play for the development of their young children. Play requires that children feel secure and that the play environment be sufficiently protected from intrusion and disruption. Parent-child play, in which the child takes the lead and the parent is attentive and responsive, elaborating but not controlling the events of play, is an excellent technique for enhancing the parent-child relationship and language development. When typical play is missing or delayed, the health care professional should consider the possibility of a developmental or emotional disorder, possible significant stresses in the child's environment, or both. The child's relationship to the family pets, if any, should be discussed and should include queries about attachment, responsibilities for pet care, and pet safety.



SEPARATION AND INDIVIDUATION

By the child's first birthday, he has likely secured a reasonably firm sense of trust that his primary caregivers are reliable, protective, and encouraging. In turn, the young toddler should begin to feel as though he can trust others enough to feel comfortable in communicating his feelings, needs, and interests. From this base of emotional security, the young child can dedicate his second year to begin growing increasingly independent from his caregivers—in actions, words, and thoughts. Periodically checking in with his parents for guidance and reassurance about safe and socially acceptable limits, the toddler waffles between testing bold new behaviors and demanding to be consoled and protected. During this stage of development, parents can help their child by providing safe opportunities for freedom and encouragement with support. As the young child develops increasing comfort in exploring time, space, and relationships with adults and peers, he begins to discover more about his own identity, effectiveness, and free will. The more positive experiences a preschool-aged

child enjoys with other children and adults, the better prepared he becomes for his subsequent adventures at school.

CHILD CARE

According to the Child Health USA 2004 Report, 63% of mothers with preschool-aged children were in the labor force (either employed or looking for employment) in 2003, with 70% of these mothers employed full-time.⁵⁵ Working mothers use a variety of child care arrangements. For example, family members are a common source of child care. A US Census Bureau report reveals²² that 40% of preschool-aged children are cared for by a relative, 23% by day care centers, nursery schools, preschools, federal Head Start programs, kindergarten, and grade schools, and 14% by family day care providers and nonrelatives (eg, babysitters, nannies, and housekeepers). Child care arrangements vary across states, ages, health status, and family income levels. Children from lower-income families are less likely to be cared for in centers than children from higher-income families, and are more likely to be in the care of relatives.⁵⁶

Families with young children, especially those living at or near the poverty level and those with several children in child care, often find that child care costs strain their budget, requiring them to balance competing family needs. Although federal subsidies for child care exist, most communities have waiting lists for openings. The health care professional and support staff who are familiar with community resources and sensitive to families' financial struggles can guide families as they make child care decisions.

Because child care for children with special health care needs is the most difficult to find and is in the shortest supply in most communities, a family's search for suitable child care can be frustrating and can sometimes cause a parent to stop working. This problem is compounded for families with low incomes,

children who have more severe special health care needs, or both. In these situations, the health care professional and staff can help families by understanding their unique needs and the available community resources. The health care professional and staff also can work with the child care provider to ensure that the setting is appropriate and the staff has the training necessary to give the child a safe and healthy environment.

Preschools should never have more than 10 children per teacher. Providers for children with special health care needs may require specialized training and support. Parents should inquire whether their preschools adhere to national standards and are accredited by organizations such as the National Association for the Education of Young Children (www.naeyc.org).²³

Quality child care gives young children valuable opportunities to learn to relate effectively with peers and adults, to explore the diverse physical and social world, and to develop confidence in their abilities to learn new skills, form trusting bonds of friendship, and process information from a variety of sources. Health care professionals should learn about the health, developmental, and behavioral issues of their patients as they are manifested in child care. Health care professionals can integrate this information in their assessment, counseling, and advocacy for children and families in their practice and their community. The more sources of insight into the child's life the health care professional has, the better prepared he will be to support the child's health and development as she takes her first steps beyond the family. Many health care professionals provide formal consultative services to child care centers in their communities.

SCHOOL READINESS

At the end of the early childhood developmental stage, the young child and his parents will begin the transition into kindergarten.

Quality child care gives young children valuable opportunities to learn to relate effectively with peers and adults, to explore the diverse physical and social world, and to develop confidence in their abilities to learn new skills, form trusting bonds of friendship, and process information from a variety of sources.

Qualities that are crucial to learning and are dependent on early emotional and social development include self-confidence, curiosity, self-control of strong emotions, motivation to learn, and the ability to make friends and become engaged in a social group.

The child will be challenged to demonstrate developmental capacities, including:

- Language and speech that is sufficient for communication and learning
- Cognitive abilities that are necessary for learning sound-letter associations, spatial relations, and number concepts
- Ability to separate from family and caregivers (especially for the child who has not already participated in preschool activities)
- Self-regulation with respect to behavior, emotions, attention, and motor movement
- Ability to make friends and get along with peers
- Ability to participate in group activities
- Ability to follow rules and directions
- Skills that others appreciate, such as singing or drawing

However, too many children today enter kindergarten significantly behind their peers in 1 or more of these abilities. Problems in self-regulation of emotions and behavior and problems in maintaining attention and focus are common at kindergarten entry and predict future educational and social problems.⁵⁷

In an extensive survey, kindergarten teachers reported that roughly half of kindergartners have difficulty following directions because of poor academic skills or problems with working in a group.⁵⁸ Racial and ethnic disparities have been shown to exist at kindergarten entry in terms of children's readiness to learn.⁵⁹

Social and emotional development during early childhood (which was neglected in past research on school readiness) has been shown to be strongly connected to later academic success. Qualities that are crucial to learning and are dependent on early emotional and social development include self-confidence, curiosity, self-control of strong emotions, motivation to learn, and the ability to make friends and become engaged in a

social group.⁵ Box 5 lists risk factors for school readiness.

The goal of having every child ready for school is a task that encompasses all of early childhood and depends on the efforts of everyone involved in the care of the young child during his first 5 years. Throughout these years, the health care professional plays a vital role in promoting this goal through assessing and monitoring the:

- General health of the child, including vision and hearing
- Child's developmental trajectory

BOX 5

Monitoring Risk Factors for School Readiness

Child-based risk factors. Readily apparent disabilities are found in a small proportion of those entering school (around 2%). Less-obvious special needs, due to specific learning disabilities, mild mental retardation, or emotional and social maladjustments, are found in nearly 20% of children.⁶⁰

Family-based risk factors. Risk factors may begin very early in childhood and include low maternal education, single-parent status, poverty, low parental literacy, households with few or no books or other reading materials, and English as a second language.⁶¹ Family habits related to television viewing also may affect school readiness. It appears that the mediators that influence readiness are the content (with children's informational shows more enhancing than general audience shows) and context (characteristics of the household) rather than television itself.

School-based risk factors. Schools may fail to recognize or accommodate the special health problems, developmental needs, and significant cultural differences of their incoming students.⁶²

DEVELOPMENTAL MILESTONES AT A GLANCE — EARLY CHILDHOOD ³⁸⁻⁶³				
Age	Gross Motor	Fine Motor	Cognitive, Linguistic, and Communication	Social-Emotional
1 Year	<ul style="list-style-type: none"> •Stand alone •Pull to stand 	<ul style="list-style-type: none"> •Put block in cup •Bang 2 cubes held in hands 	<ul style="list-style-type: none"> •Imitate vocalizations and sounds •Babbling* •1 word 	<ul style="list-style-type: none"> •Protodeclarative pointing* •Wave bye-bye •Imitate activities •Play pat-a-cake
15 Months	<ul style="list-style-type: none"> •Walk backwards •Stoop and recover •Walk well 	<ul style="list-style-type: none"> •Scribble •Put block in cup 	<ul style="list-style-type: none"> •1 word* •3 words 	<ul style="list-style-type: none"> •Drink from cup •Wave bye-bye
18 Months	<ul style="list-style-type: none"> •Walk up steps •Run •Walk backwards 	<ul style="list-style-type: none"> •Dump raisin, demonstrated •Tower of 2 cubes •Scribble 	<ul style="list-style-type: none"> •Point to at least 1 body part •6 words •3 words 	<ul style="list-style-type: none"> •Remove garment •Help in house
2 Years	<ul style="list-style-type: none"> •Throw ball overhand •Jump up •Kick ball forward •Walk up steps 	<ul style="list-style-type: none"> •Tower of 6 cubes •Tower of 4 cubes 	<ul style="list-style-type: none"> •Name 1 picture •Combine words •Point to 2 pictures 	<ul style="list-style-type: none"> •Put on clothing •Remove garment
2½ Years	<ul style="list-style-type: none"> •Throw ball overhand •Jump up 	<ul style="list-style-type: none"> •Imitate vertical line •Tower of 8 cubes •Tower of 6 cubes 	<ul style="list-style-type: none"> •Know 2 actions •Speech half understandable •Point to 6 body parts •Name 1 picture 	<ul style="list-style-type: none"> •Wash and dry hands •Put on clothing
3 Years	<ul style="list-style-type: none"> •Balance on each foot 1 second •Broad jump •Throw ball overhand 	<ul style="list-style-type: none"> •Thumb wiggle •Imitate vertical line •Tower of 8 cubes •Tower of 6 cubes 	<ul style="list-style-type: none"> •Speech all understandable •Name 1 color •Know 2 adjectives •Name 4 pictures 	<ul style="list-style-type: none"> •Name friend •Brush teeth with help
4 Years	<ul style="list-style-type: none"> •Hop •Balance on each foot 2 seconds 	<ul style="list-style-type: none"> •Draw a person with 3 parts •Tower of 8 cubes 	<ul style="list-style-type: none"> •Define 5 words •Name 4 colors •Speech all understandable 	<ul style="list-style-type: none"> •Copy a cross (+) •Copy a circle

KEY

Black Color: 50% to 90% of children pass this item.

Green Color: More than 90% of children pass this item.

***Absence of these milestones should trigger screening for autism.**

These norms are taken from the DENVER II, and are based upon the administration and interpretation as set forth in the DENVER II Training Manual (copyright 1992).

These milestones are provided as a reference only. Reference to these milestones does not take the place of a standardized measurement of healthy child development or discourage a developmental discussion with a health care provider.

Middle childhood is an important time for families to help children consolidate and strengthen their cognitive and emotional attributes, such as communication skills, sensitivity to others, ability to form positive peer relationships, self-esteem, and independence.

- Emotional health of the child and family, especially when based on the health care professional's long-term knowledge of child-family relationships
- Child's social development (both skills and difficulties)
- Specific child-based, family-based, school-based, and community-based risk factors

Health care professionals have a unique opportunity to recognize problems and, when possible, to intervene early with effective referral for both specific services and general evaluation so as to enhance the child's readiness for learning by the start of school. Intervention services for eligible children can begin at birth and continue through age 21 years. For details on eligibility and services, refer to the US Department of Education's Office of Special Education and Rehabilitative Services (<http://www.ed.gov/about/offices/list/osers>).

Promoting Child Development: Middle Childhood—5 to 10 Years

The middle childhood years are typically a stable period, without the dramatic physical, cognitive, and social changes that occur in the other developmental stages. It is an important transitional period during which children build on the skills developed in the various domains of early childhood in preparation for adolescence. Middle childhood is an important time for families to help

children consolidate and strengthen their cognitive and emotional attributes, such as communication skills, sensitivity to others, ability to form positive peer relationships, self-esteem, and independence. These attributes will help them cope with the stresses and potential risks of adolescence.

Families and health care professionals also can provide enormous support for children's healthy physical development. They can work with communities to ensure that children have access to safe, well-supervised play areas, recreation centers, team sports and organized activities, parks, and schools. For children to flourish, communities must provide carefully maintained facilities to help their bodies and minds develop in a healthy way. Health care professionals can support their guidance by advocating for community facilities available to all children. (For more information on this topic, see the Promoting Community Relationships and Resources and the Promoting Physical Activity themes.)

Children and Youth With Special Health Care Needs

Middle childhood is a critical time for children with special health care needs to adapt successfully to their condition. During this period, they continue to define their sense of self and improve their ability to care for their own health. Children adapt best to chronic illness when health care professionals, families, schools, and communities work together to foster their emerging independence. Inclusion in school and community life allows children with special health care needs to feel valued and to integrate their specific care needs with other aspects of their lives. It also is important to discuss family perspectives, because families may have various beliefs and values regarding the independence of children with special health care needs based on culture and history.

When families have children with special health care needs, the other children in the



family will be introduced to the possibility that their sibling can have different challenges because of the disability and circumstances. Families may have to deal with certain difficult tasks, such as hospitalizations or painful tests, illness, and possibly death. Parents and child care providers should be sensitive to these issues and responsive to the needs of the medically fragile child and the healthy siblings. At the same time, children with special health care needs should not be given special privileges simply because of their condition. Instead, outlining rules and responsibilities is extremely important for the child's development and the family's functioning. Child care providers and teachers can play an important supportive role and be a source of information for the parents and the children.

Domains of Development

GROSS AND FINE MOTOR SKILLS

Major increases in strength and improvements in motor coordination occur during middle childhood. These changes contribute to the child's growing sense of competence in relation to his physical abilities and enhance his potential for participating in sports, dance, gymnastics, and other physical pursuits. A child's participation in sports or other physical activities can reinforce positive interaction skills that will serve the child throughout his life. Children develop at slightly different rates depending on their unique physical characteristics and experiences. Efforts to maintain good physical health and exercise patterns are important to achieving and maintaining a healthy weight. Monitoring the child's growth patterns and conducting periodic physical examinations to assess growth and development are important components of health supervision. Close monitoring for overweight and obesity should be included in these visits. (For more information on this topic, see the Promoting Healthy Weight theme.)

COGNITIVE, LINGUISTIC, AND COMMUNICATION SKILLS

Children's readiness to learn in school depends on cognitive maturation as well as their individual experiences. During middle childhood, the child moves from magical thinking to more logical thought processes. The synthesis of basic language, perception, and abstraction allows the child to read, write, and communicate thoughts of increasing complexity and creativity. Progress can appear subtle from month to month, but it is dramatic from one school year to the next. As the child's cognitive skills grow, she matures in her ability to understand the world and people around her and to function independently. Occasionally, children will be impaired in their development because of learning problems, behavior problems, or both. The health care professional can offer support by ensuring screening and evaluation for any suspected delays.

The major developmental achievement of this age is self-efficacy, or the knowledge of what to do and the confidence and ability to do it. Success at school is most likely to occur when this achievement is encouraged by parents and valued by families. Families who reward children with enthusiasm and warmth for putting forth their best effort ensure their steady educational progress and prepare them to use their intelligence and knowledge productively. Through awareness of individual learning styles, including the need for necessary accommodations, parents and teachers can adapt materials and experiences to each child. School success is an important factor in the development of a child's self-esteem. In families in which parents have had unsuccessful educational experiences or have had limited education, support from health care professionals and others in the community is critical in supporting their children through the educational process.

During middle childhood, the child moves from magical thinking to more logical thought processes.

He becomes increasingly aware that an individual's life fits into a larger scheme of relationships among individuals, groups of people, other living creatures, and the earth itself.

SOCIAL-EMOTIONAL SKILLS

As children become increasingly independent and demonstrate initiative, they develop their own sense of personhood (Table 4). They begin to discern where they “fit” among their peers and in their family, school class, neighborhood, and community. When the “fit” is good and comfortable, children see themselves as effective and competent members of their family, group, team, school, and community. When the “fit” is tenuous or poor, the dissonance can be a source of distress and can predispose children to emotional disorders with long-term consequences. (For more information on this topic, see the [Promoting Mental Health](#) theme.) Ongoing support for the child provides the best opportunity for acceptance and forms the basis for a strong self-worth. Support is especially important for children with special health care needs.

Children need both the freedom of personal expression and the structure of expectations and guidelines that they can understand and accept. Families should provide opportunities for the child to interact with other children in play environments without excessive adult interference. However, not all cultures accept this perspective. The health care professional and the family should discuss these issues. Most experts believe that children benefit from the experience of independent play with peers. Unfortunately, some neighborhoods or living arrangements restrict these opportunities. In addition, some children with special health care needs may need adaptive equipment or facilities to allow for inclusive play experiences. Children also need to have positive interactions with adults, reinforcing their sense of self-esteem, self-worth, and belief in their capability of personal success.

The child’s “self” evolves in a social context. Health care professionals can help families understand this dynamic and encourage specific roles for the children within the

family. Parents who consciously assess their child’s emotional maturity and role in the family at each birthday will appreciate the changes that have occurred subtly over time.

Children with special health care needs will have emotional maturity that is appropriately reflective of their needs, developmental level, and physical challenges (Table 4). Parents should be encouraged to appreciate the individual maturity level of their child. As a result, they can celebrate the child’s evolving autonomy by granting new privileges. Parents who match each new entitlement with a new responsibility signal their respect for the child’s growing capability to contribute to the family and the community.

Developmental Highlights of Middle Childhood

MORAL AND SPIRITUAL DEVELOPMENT

The child’s development as an individual involves an understanding of the life cycle—birth, growth and maturation, aging, and death. He becomes increasingly aware that an individual’s life fits into a larger scheme of relationships among individuals, groups of people, other living creatures, and the earth itself. School-aged children become keenly interested in these topics, especially if they experience life events such as the birth of a sibling or the death of a grandparent. Children also become aware of violent death, on the highways or on street corners. When a death occurs, parents should be encouraged to discuss the loss with their children and provide assistance to children who are having difficulty with the grieving process.

As children experience these events and learn to view their personal encounters as part of a larger whole, families and communities provide an important structure. These experiences provide children with a basic foundation of value systems and encourage them to examine their personal actions in the context of those around them.

TABLE 4

Social and Emotional Development in Middle Childhood

Topics	Key Areas
Self	<p data-bbox="380 348 1214 380"><i>(Key areas in italics are especially important for children with special health care needs.)</i></p> <p data-bbox="380 386 529 417">Self-esteem:</p> <ul data-bbox="412 422 928 562" style="list-style-type: none"> <li data-bbox="412 422 716 453">• Experiences of success <li data-bbox="412 457 829 489">• Reasonable risk-taking behavior <li data-bbox="412 493 899 525">• Resilience and ability to handle failure <li data-bbox="412 529 928 562">• Supportive family and peer relationships <p data-bbox="380 579 516 611">Self-image:</p> <ul data-bbox="412 615 1192 751" style="list-style-type: none"> <li data-bbox="412 615 1008 646">• Body image, <i>celebrating different body images</i> <li data-bbox="412 651 1192 751">• Prepubertal changes; initiating discussion about sexuality and reproduction; <i>prepubertal changes related to physical care issues</i>
Family	<p data-bbox="380 762 667 793">What matters at home:</p> <ul data-bbox="412 798 789 1041" style="list-style-type: none"> <li data-bbox="412 798 789 829">• Expectation and limit setting <li data-bbox="412 833 708 865">• Family times together <li data-bbox="412 869 643 900">• Communication <li data-bbox="412 905 703 936">• Family responsibilities <li data-bbox="412 940 654 972">• Family transitions <li data-bbox="412 976 686 1008">• Sibling relationships <li data-bbox="412 1012 724 1041">• <i>Caregiver relationships</i>
Friends	<p data-bbox="380 1052 521 1083">Friendships:</p> <ul data-bbox="412 1087 1214 1220" style="list-style-type: none"> <li data-bbox="412 1087 1214 1150">• Making friends, <i>friendships with peers with and without special health care needs</i> <li data-bbox="412 1155 1175 1220">• Family support of friendships, <i>family support to have typical friendship activities, as appropriate</i>
School	<p data-bbox="380 1230 472 1262">School:</p> <ul data-bbox="412 1266 1203 1577" style="list-style-type: none"> <li data-bbox="412 1266 1138 1329">• Expectation for school performance, <i>school performance defined in the Individualized Education Program (IEP)</i> <li data-bbox="412 1333 581 1365">• Homework <li data-bbox="412 1369 1154 1400">• Child-teacher conflicts, <i>building relationships with teachers</i> <li data-bbox="412 1404 816 1436">• <i>Parent-teacher communication</i> <li data-bbox="412 1440 1203 1503">• Ability of schools to address the needs of children from diverse backgrounds <li data-bbox="412 1507 1073 1539">• Awareness of aggression, bullying, and victimization <li data-bbox="412 1543 602 1577">• Absenteeism
Community	<p data-bbox="380 1587 651 1619">Community strengths:</p> <ul data-bbox="412 1623 760 1724" style="list-style-type: none"> <li data-bbox="412 1623 760 1654">• Community organizations <li data-bbox="412 1659 646 1690">• Religious groups <li data-bbox="412 1694 634 1724">• Cultural groups <p data-bbox="380 1728 846 1759">High-risk behaviors and environments:</p> <ul data-bbox="412 1764 1214 1932" style="list-style-type: none"> <li data-bbox="412 1764 618 1795">• Substance use <li data-bbox="412 1799 667 1831">• Unsafe friendships <li data-bbox="412 1835 846 1866">• Unsafe community environments <li data-bbox="412 1871 1214 1932">• <i>Particular awareness of risk-taking behaviors and unsafe environments, because children may be easily victimized</i>

As children with special health care needs enter adolescence and experience puberty, growth, and physical and emotional development, new levels of functionality in the face of their special need can bring important and remarkable gains in independence and autonomy.

The relationship between values, competence, self-esteem, and personal responsibility needs to be modeled and affirmed by the child's parents, teachers, and communities. Parents need to help their child maintain a balance of responsibilities at school and home, time spent with family and friends, extracurricular and community activities, and personal leisure. Achieving this balance is essential for healthy development. Competence and self-esteem are strengthened when a child is recognized for working hard in school, successfully completing chores and special projects, and participating in school and community activities.

Promoting Child Development: Adolescence—11 to 21 Years

Adolescence is a dynamic experience, not a homogenous period of life. Adolescents differ widely in their physical, social, and emotional maturity because they enter puberty at different ages, progress at different paces, and experience different challenges in their developmental trajectories. To complicate the adolescent experience, parents also can experience changes in health, employment, geographic relocation, marital relationships, or the health of their parents and other family members.

Viewing adolescence in stages—early adolescence (11 to 14 years of age), middle adolescence (15 to 17 years of age), and late adolescence (18 to 21 years of age)—yields a better understanding of physical and psychological development and potential problems. The nature, length, and course of typical adolescent development can be viewed differently by families because cultural expectations for independence and self-sufficiency can differ. The health care professional should discern from families how they view this stage of life and note potential conflicts between the family's cultural values and those of the developing adolescent.

Youth With Special Health Care Needs

As children with special health care needs enter adolescence and experience puberty, growth, and physical and emotional development, new levels of functionality in the face of their special need can bring important and remarkable gains in independence and autonomy. Alternatively, limitations related to their illness can further underscore their physical dependence, which can limit the development of emotional independence. The adolescent may fear that his condition precludes autonomy.

Careful assessment of medical conditions, strengths and risk-taking behaviors, followed by sensitive discussions of the youth's perceived needs and goals, can assist the adolescent with a special health care need to maximize physical development and support the attainment of full emotional development and maturity.

Stages of Adolescence

Three key transitional domains (physiological, psychological, and social) can be used to chart adolescent changes and challenges (Table 5).

Domains of Development

GROSS AND FINE MOTOR SKILLS

Pubertal growth brings completion of physical development. Adult height and muscle mass are attained. Increasing size and



TABLE 5

Domains of Adolescent Development

	Early Adolescence (11 to 14 Years)	Middle Adolescence (15 to 17 Years)	Late Adolescence (18 to 21 Years)
Physiological	Onset of puberty, growth spurt, menarche (females)	Ovulation (females), growth spurt (males)	Growth completed
Psychological	Concrete thought, preoccupation with rapid body changes, sexual identity, questioning independence, parental controls remain strong	Competence in abstract and future thought, idealism, sense of invincibility or narcissism, sexual identity, beginning of cognitive capacity to provide legal consent	Future orientation, emotional independence, unmasking of psychiatric disorders, capacity for empathy, intimacy, and reciprocity in interpersonal relationships, self-identity, recognized as legally capable of providing consent, ⁶⁴ attainment of legal age for some issues (eg, voting) but not all issues (eg, drinking alcohol)
Social	Search for same-sex peer affiliation, good parental relationships, other adults as role models; transition to middle school, involvement in extracurricular activities; sensitivity to differences between home culture and culture of others	Beginning emotional emancipation, increased power of peer group, conflicts over parental control, interest in sexual relationships, initiation of driving, risk-taking behavior, transition to high school, reduced involvement in extracurricular activities, possible cultural conflict as adolescent navigates between family's values and values of broader culture and peer culture	Individual over peer relationships; transition in parent-child relationship, transition out of home; may begin preparation for further education, career, marriage, and parenting
Potential Problems	Delayed puberty; acne; orthopedic problems; school problems; psychosomatic concerns; depression; unintended pregnancy; initiation of tobacco, alcohol, or other drug use	Experimentation with health risk behaviors (eg, sex, drinking, drug use, smoking), auto crashes, menstrual disorders, unintended pregnancy, acne, short stature (males), conflicts with parents, overweight, physical inactivity, poor eating behaviors, eating disorders (eg, purging, binge eating, and anorexia nervosa)	Eating disorders, depression, suicide, auto crashes, unintended pregnancy, acne, smoking, alcohol or drug dependence

Parental involvement and expectations and participation in extracurricular activities enhance adolescent academic achievement and educational attainment.

strength are accompanied by enhanced coordination of both gross and fine motor skills. The boy or girl who can barely make the high-school junior varsity basketball team as a ninth grader has the agility and strength necessary for varsity performance by 10th or 11th grade. Motor development continues into the final stage of development.

COGNITIVE, LINGUISTIC, AND COMMUNICATION SKILLS

Success in school contributes substantially to the adolescent's self-esteem and progress toward becoming a socially competent adult. The National Longitudinal Study for Adolescent Health⁶⁵ found that school performance and choice of free-time activities were the most important determinants for every risky behavior studied, regardless of socioeconomic status, race, or type of household (ie, 1 parent vs 2 parents). Students who have a high academic self-concept tend to be more motivated to achieve, more engaged in school, and more hopeful about their future. Parental involvement and expectations and participation in extracurricular activities enhance adolescent academic achievement and educational attainment. Health care professionals should encourage conversations between parents and their adolescent children on these issues.

Adolescents who feel connected to their school and who have a high academic self-concept are motivated to achieve. Peer relationships also influence adolescents' attitudes. Adolescents whose peers have or are perceived to have higher educational aspirations tend to be more engaged in school and to have higher hopes for continuing their education. Conversely, adolescents who work more than 20 hours a week tend to have a lower level of engagement in school.⁶⁶ The health care professional should encourage youth to participate in extracurricular activities. Factors such as disability and limited English proficiency can interfere with school success and need attention.

Some adolescents make the academic and social transition from middle school to high school easily. Others find this transition overwhelming, with an impact on motivation, self-esteem, and academic performance.

The Centers for Disease Control and Prevention (CDC) National Center for Health Statistics estimates that among children 3 to 17 years of age, 8% have a learning disability.⁶⁷ Children with a fair or poor health status were 5 times as likely to have a learning disability than children with an excellent or a very good health status.⁶⁷ Students with learning disabilities can have difficulty with academics as well as social relationships. These students are more prone to depression and a lack of confidence.⁶⁸ Health care professionals should screen youth for declining grades and attendance issues, signs of learning disorders, and social adjustment concerns. Depending on the specific school district policies, health care professionals can interact with the school nurse, psychologist, counselor, or administrator to identify and address academic, social, and emotional difficulties that can interfere with school success.

SOCIAL-EMOTIONAL SKILLS

A consistent, supportive environment for the adolescent, with graded steps toward autonomy is necessary to foster emotional and social well-being. This supportive environment requires the participation of the family, school, health care professional, and community, as well as the adolescent himself.⁶⁹ Parents will struggle for a balance for their child between restrictions that are designed to protect him and freedom that is intended to enhance growth. The adolescent will struggle for this balance, too.

The emotional well-being of adolescents is tied to their sense of self-esteem. High self-esteem is generally associated with feelings of life satisfaction and a sense of control over one's life, whereas low self-esteem is correlated with lower reports of happiness and

higher reports of feeling as if one is not in control of one’s life. Adolescents who demonstrate good social and problem-solving skills also usually have enhanced self-esteem because these skills increase their sense of control over their world. This asset is essential in deriving the ability to handle stress and cope with challenging situations.

Another important developmental milestone that is critical to emotional well-being is the adolescent’s growing sense of self. Long hours spent talking, grooming, being alone, and rushing to be part of a group—any group—are all part of the adolescent’s search for a conception of self. Intelligence, in the narrow sense of the term, also is significant to the cognitive self. During adolescence, the individual has to learn the accumulated wisdom of society. As the adolescent becomes facile in using concepts and abstractions, he begins to combine new ideas in new ways to arrive at creative solutions.

Normal fluctuations of mood now are the adolescent’s responsibility. With increasing autonomy, he may become unwilling to share feelings and, to a point, unconsciously seek to avoid dependence on family for mood modulation. Like other skills he acquires, managing feelings of sadness and anxiety requires guidance, practice, and experience.

During the course of adolescence, the increasingly autonomous and socially competent youth finds his place in family and community. Social competence is defined as the degree to which significant others rate an individual as successful in solving and completing social tasks.⁷⁰ The specific behaviors that characterize social competence will vary with the situation in which the adolescent is functioning.⁷¹ Socially competent youth are able to decode and interpret social cues and consider alternative responses, along with their consequences.

To function in an adult world, a youth must become aware of his relations to others and learn the personal impact of relationships

on his daily function. Accordingly, he must appreciate the effects and impact of his actions toward others if relationships are to be mature and reciprocal. Understanding how others might interpret a situation, recognizing another’s predicament, and comfortably appreciating another’s feelings are new and important experiences. Empathy must be achieved for healthy adult relationships to flourish.

The adolescent’s social and emotional skills also are influenced by the young adult’s growing interactions with the wider community through travel, higher education, volunteer activities, or structured job experiences. These activities can help adolescents realize that they have meaningful roles and can contribute productively to society. Through these activities, youth learn the importance of general adherence to rules and authority. External mandates are internalized in an appreciation of right or wrong and consequences.

Developmental Highlights of Adolescence

ASSETS

Health advocates have begun to look at the family and community factors that promote healthy development. This “asset” model, or strength-based approach, provides a broader perspective on adolescent development than the more traditional “deficit” model, which looks at the problems experienced by adolescents and develops preventive interventions (Table 6). The asset model reinforces

TABLE 6

Comparison of Asset and Deficit Models

Asset Model	Deficit Model
Positive family environment	Abuse or neglect
Relationships with caring adults	Witness to domestic violence
Religious and spiritual anchors	Family discord and divorce
Involvement in school, faith-based organization, or community	Parents with poor health habits
Accessible recreational opportunities	Unsafe schools
	Unsafe neighborhood

When young people decide to seek assistance beyond their family, those resources should provide appropriate confidential counseling and support to them in making healthy choices while encouraging good communication with parents and family.

health-promoting interactions or social involvement (eg, good parent-adolescent communication and participation in extracurricular activities) and assists adolescents and their parents in setting goals to achieve healthy development.

Research demonstrates the value of parental involvement and quality parent-adolescent communication on healthy adolescent development.⁷²⁻⁷⁴ Adolescents whose parents are authoritative, rather than authoritarian or passive, and who are involved in extracurricular and community activities appear to progress through adolescence with relatively little turmoil.⁷⁵

MODELS OF CARE

Onsite integrated health services in the schools—with referrals to health care professionals and community agencies (including mental health centers) for supplementary services—are evolving as one way to deliver adolescent health care in medically underserved areas. In some situations, the school-based health center is the medical home for the youth enrolled in the center. School-based health centers can be especially effective in ensuring immunizations, promoting sports safety, and providing access for students with special health care needs. All services and programs should work to improve communication between school and home so that parents stay involved in their adolescents' lives away from home and learn effective strategies to deal with some of the challenges that their children face.

Health care professionals should ask young people how they learn about healthy living. Health promotion programs in schools help adolescents establish good health habits and avoid those that can lead to morbidity and mortality. Health promotion curricula can include family life education and social skills training, as well as information on pregnancy prevention, abstinence, conflict resolution, healthy nutritional practices, and avoidance of unhealthy habits such as the use of tobacco products, alcohol, or other drugs. Referrals to appropriate, culturally respectful, and accessible community resources also help adolescents learn about and address mental health concerns, nutrition and physical health, and sexual health issues. When young people decide to seek assistance beyond their family, those resources should provide appropriate confidential counseling and support to them in making healthy choices while encouraging good communication with parents and family.



References

1. Council on Children with Disabilities; Section on Developmental Behavioral Pediatrics; Bright Futures Steering Committee; Medical Home Initiatives for Children with Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening. *Pediatrics*. 2006;118:405-420: Erratum in: *Pediatrics*. 2006;188:1808-1809
2. Glascoe FP, Dworkin PH. The role of parents in the detection of developmental and behavioral problems. *Pediatrics*. 1995;95:829-836
3. Jellinek MS, Patel BP, Froehle MC, eds. *Bright Futures in Practice: Mental Health. Tool Kit—Volume 2*. Arlington, VA: National Center for Education in Maternal and Child Health; 2002
4. Knight JR, Sherritt L, Shrier LA, Harris SK, Chang G. Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. *Arch Pediatr Adolesc Med*. 2002;156:607-614
5. National Research Council and Institute of Medicine, Committee on Integrating the Science of Early Childhood Development. *From Neurons to Neighborhoods: The Science of Early Childhood Development*. Shonkoff JP, Phillips DA, eds. Washington, DC: National Academy Press; 2000
6. Dodd VL. Implications of kangaroo care for growth and development in preterm infants. *J Obstet Gynecol Neonatal Nurs*. 2005;34:218-232
7. Feldman R, Eidelman AI, Sirota L, Weller A. Comparison of skin-to-skin (kangaroo) and traditional care: parenting outcomes and preterm infant development. *Pediatrics*. 2002;110:16-26
8. Shackelford J. State and jurisdictional eligibility definitions for infants and toddlers with disabilities under IDEA. NECTAC Notes No. 21. Chapel Hill: The University of North Carolina, FPG Child Development Institute, National Early Childhood Technical Assistance Center; 2006. Available at: <http://www.nectac.org/%7Epdfs/pubs/nnotes21.pdf>. Accessed March 1, 2006
9. Shore R. What have we learned? In: *Rethinking the Brain: New Insights into Early Development*. New York, NY: Families and Work Institute; 1997:15-55
10. Weinberger DR, Elvevag B, Giedd JN. *The Adolescent Brain: A Work in Progress*. Washington, DC: The National Campaign to Prevent Teen Pregnancy; 2005
11. American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. The changing concept of sudden infant death syndrome: diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. *Pediatrics*. 2005;116:1245-1255
12. Gorski PA. Contemporary pediatric practice: in support of infant mental health (imaging and imagining). *Infant Ment Health J*. 2001;22:188-200
13. American Academy of Pediatrics, Committee on Public Education. Children, adolescents, and television. *Pediatrics*. 2001;107:423-426
14. Wright JC, Huston AC, Murphy KC, et al. The relations of early television viewing to school readiness and vocabulary of children from low-income families: the early window project. *Child Dev*. 2001;72:1347-1366
15. Atkinson PM, Parks DK, Cooley SM, Sarkis SL. Reach Out and Read: a pediatric clinic-based approach to early literacy promotion. *J Pediatr Health Care*. 2002;16:10-15
16. High PC, LaGasse L, Becker S, Ahlgren I, Gardner A. Literacy promotion in primary care pediatrics: can we make a difference? *Pediatrics*. 2000;105:927-934
17. Sharif I, Rieber S, Ozuah PO. Exposure to Reach Out and Read and vocabulary outcomes in inner city preschoolers [published erratum appears in *J Natl Med Assoc*. 2002;94(9)]. *J Natl Med Assoc*. 2002;94:171-177
18. Weitzman CC, Roy L, Walls T, Tomlin R. More evidence for reach out and read: a home-based study. *Pediatrics*. 2004;113:1248-1253
19. Mendelsohn AL, Mogilner LN, Dreyer BP, et al. The impact of a clinic-based literacy intervention on language development in inner-city preschool children. *Pediatrics*. 2001;107:130-134
20. Needlman R, Silverstein M. Pediatric interventions to support reading aloud: how good is the evidence? *J Dev Behav Pediatr*. 2004;25:352-363
21. Needlman R, Toker KH, Dreyer BP, Klass P, Mendelsohn AL. Effectiveness of a primary care intervention to support reading aloud: a multicenter evaluation. *Ambul Pediatr*. 2005;5:209-215
22. Overturf Johnson JO. *Who's Minding the Kids? Child Care Arrangements: Winter 2002*. Washington, DC: US Census Bureau; 2005 (Current Population Reports. Household Economic Studies p70, 101)
23. American Academy of Pediatrics, Committee on Early Childhood, Adoption, and Dependent Care. Quality early education and child care from birth to kindergarten. *Pediatrics*. 2005;115:187-191
24. Carlson VJ, Harwood RL. What do we expect? Understanding and negotiating cultural differences concerning early developmental competence: the six raisin solution. *Zero to Three*. 1999;20:19-24
25. National SIDS/Infant Death Resource Center. What is SIDS? Available at: <http://www.sidscenter.org/WhatsSIDS.pdf>. Accessed September 4, 2007
26. Barnard KE. *Beginning Rhythms: The Emerging Process of Sleep Wake Behaviors and Self-Regulation*. Seattle, WA: NCAST, University of Washington; 1999

27. Carno MA, Hoffman LA, Carcillo JA, Sanders MH. Developmental stages of sleep from birth to adolescence, common childhood sleep disorders: overview and nursing implications. *J Pediatr Nurs*. 2003;18:274-283
28. Scheers NJ, Rutherford GW, Kemp JS. Where should infants sleep? A comparison of risk for suffocation of infants sleeping in cribs, adult beds, and other sleeping locations. *Pediatrics*. 2003;112:883-889
29. Bornstein MH. *Cultural Approaches to Parenting*. Hillsdale, NJ: Lawrence Erlbaum; 1991
30. Harkness S, Super CM, Keefer CH, van Tuijn N, van der Klugt E. *Cultural Influences on Sleep Patterns in Infancy and Early Childhood*. Atlanta, GA: American Association for the Advancement of Science; 1995
31. Brenner RA, Simons-Morton BG, Bhaskar B, Revenis M, Das A, Clemens JD. Infant-parent bed sharing in an inner-city population. *Arch Pediatr Adolesc Med*. 2003;157:33-39
32. Nakamura S, Wind M, Danello MA. Review of hazards associated with children placed in adult beds. *Arch Pediatr Adolesc Med*. 1999;153:1019-1023
33. Unger B, Kemp JS, Wilkins D, et al. Racial disparity and modifiable risk factors among infants dying suddenly and unexpectedly. *Pediatrics*. 2003;111(2):e127-e131. Available at: www.pediatrics.org/cgi/content/full/111/2/e127. Accessed May 4, 2007
34. Willinger M, Ko CW, Hoffman HJ, Kessler RC, Corwin MJ. Factors associated with caregivers' choice of infant sleep position, 1994-1998: the National Infant Sleep Position Study. *JAMA*. 2000;283:2135-2142
35. Li DK, Petitti DB, Willinger M, et al. Infant sleeping position and the risk of sudden infant death syndrome in California, 1997-2000. *Am J Epidemiol*. 2003;157:446-455
36. Sivan Y, Reisner S, Amitai Y, Wasser J, Nehama H, Tauman R. Effect of religious observance on infants' sleep position in the Jewish population. *J Paediatr Child Health*. 2004;40:534-539
37. Willinger M, Ko CW, Hoffman HJ, Kessler RC, Corwin MJ. Trends in infant bed sharing in the United States, 1993-2000: the National Infant Sleep Position study. *Arch Pediatr Adolesc Med*. 2003;157:43-49
38. Frankenburg WK, Dodds J, Archer P, et al: DENVER II Training Manual, 1992. Denver Developmental Materials. Denver, CO
39. Diego MA, Field T, Hart S, et al. Facial expressions and EEG in infants of intrusive and withdrawn mothers with depressive symptoms. *Depress Anxiety*. 2002;15:10-17
40. Diego MA, Field T, Hernandez-Reif M, Cullen C, Schanberg S, Kuhn C. Prepartum, postpartum, and chronic depression effects on newborns. *Psychiatry*. 2004;67:63-80
41. Field T. Early interventions for infants of depressed mothers. *Pediatrics*. 1998;102(5 Suppl E):1305-1310
42. Bowman BT. Educating language minority children: challenges and opportunities. *Phi Delta Kappan*. 1989;71:118-120
43. Kopp CB. Antecedents of self-regulation: a developmental perspective. *Dev Psychol*. 1982;18:199-214.
44. Raffaelli M, Crockett LJ, Shen YL. Developmental stability and change in self-regulation from childhood to adolescence. *J Genet Psychol*. 2005;166:54-75
45. Thiedke CC. Nocturnal enuresis. *Am Fam Physician*. 2003;67:1499-1506
46. Evans GD. *Bedwetting*. Gainesville, FL: Department of Family, Youth, and Community Sciences, Florida Cooperative Extension Service, Institute of Food and Agriculture Sciences, University of Florida; 2003. Available at: <http://edis.fas.ufl.edu/pdffiles/HE/HE79400.pdf>. Accessed September 13, 2007
47. Search Institute. *Developmental Assets*. Minneapolis, MN: Search Institute; 1997. Available at: <http://www.search-institute.org/assets>. Accessed September 5, 2007
48. Scales P, Leffert N, Lerner RM. *Developmental Assets: A Synthesis of the Scientific Research on Adolescent Development*. Minneapolis, MN: Search Institute; 1999
49. Biles B. Activities that promote racial and cultural awareness. *Family Child-Care Connections*. 1994;4:1-4. Available at: http://web.aces.uiuc.edu/vista/pdf_pubs/CHILDCARE.PDF. Accessed September 13, 2007
50. American Academy of Pediatrics, Committee on Psychosocial Aspects of Child and Family Health. Guidance for effective discipline. *Pediatrics*. 1998;101:723-728. Erratum in: *Pediatrics*. 1998;102:433
51. Burns MS, Griffin P, Snow CE. *Starting Out Right: A Guide to Promoting Children's Reading Success*. Washington, DC: National Academy Press; 1999
52. Podhajski B, Nathan J. Promoting early literacy through professional development for childcare providers. *Early Educ Dev*. 2005;16:1-5
53. Zuckerman BS, Klass P. Doctors Promoting Child Development with Books. Available at: http://www.ced.org/docs/report/report_ivk_zuckerman_2006.pdf. Accessed September 5, 2007
54. Glassy D, Romano J, American Academy of Pediatrics, Committee on Early Childhood, Adoption, and Dependent Care. Selecting appropriate toys for young children: the pediatrician's role. *Pediatrics*. 2003;111:911-913
55. US Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau. *Child Health USA 2004*. Rockville, MD: US Department of Health and Human Services; 2004. Available at www.mchb.hrsa.gov/mchirc/chusa_04/pdf/eo4.pdf. Accessed September 13, 2007
56. Capizzano J, Adams G, Sonenstein F. *Child Care Arrangements for Children Under Five: Variation Across States*. Washington, DC: Urban Institute; 2000
57. Raver CC. *Emotions Matter: Making the Case for the Role of Young Children's Emotional Development for Early School Readiness*. Ann Arbor, MI: Society for Research in Child Development; 2002
58. Rimm-Kaufman SE, Pianta RC, Cox MJ. Teachers' judgments of problems in the transition to kindergarten. *Early Child Res Q*. 2000;15:147-166

59. Rouse C, Brooks-Gunn J, McLanahan S. School readiness: closing racial and ethnic gaps: introducing the issue. *Future Child*. 2005;15:5-14
60. Farran DC, Shonkoff JP. Developmental disabilities and the concept of school readiness. *Early Educ Dev*. 1994;5:141-151
61. West J, Flanagan KD, Germino-Hausken E. *America's Kindergartners: Findings from the Early Childhood Longitudinal Study, Kindergarten Class of 1998-99 Fall 1998*. Washington, DC: US Department of Education, National Center for Education Statistics; 1998. Publication No. NCES 2000070
62. Meisels SJ. Assessing readiness. In: Pianta RC, Cox MJ, eds. *The Transition to Kindergarten*. Baltimore, MD: Paul H Brookes; 1999:39-66
63. Coplan J. *Early Language Milestone Scale*. 2nd ed. Austin, TX: Pro-Ed Inc; 1993
64. English A, Kenney KE. *State Minor Consent Laws: A Summary*. 2nd Ed. Chapel Hill, NC: Center for Adolescent Health & The Law; 2003
65. Resnick MD, Bearman PS, Blum RW, et al. Protecting adolescents from harm. Findings from the National Longitudinal Study on Adolescent Health. *JAMA*. 1997;278:823-832
66. Redd Z, Brooks J, McGarvey AM. *Educating America's Youth: What Makes a Difference*. Washington, DC: Child Trends; 2002
67. Bloom B, Dey AN. Summary health statistics for U.S. children: National Health Interview Survey, 2004. *Vital Health Stat 10*. 2006;Feb(227):1-85
68. US Department of Education. *Helping Your Child Through Early Adolescence for Parents of Children from 10 through 14*. Washington, DC: US Department of Education. Available at: <http://www.ed.gov/parents/academic/help/adolescence/adolescence.pdf>. Accessed September 5, 2007
69. Irwin CE. The adolescent, health, and society: from the perspective of the physician. In: Millstein SG, Peterson AC, Nightingale EO, eds. *Promoting the Health of Adolescents: New Directions for the Twenty-first Century*. New York, NY: Oxford University Press; 1993:146-150
70. Dodge KA. A social information processing model of social competence in children. In: Perlmutter M, ed. *Minnesota Symposium on Child Psychology*. Vol 18. Hillsdale, NJ: Lawrence Erlbaum Associates; 1986:77-125
71. Hair EC, Jager J, Garrett SB. *Helping Teens Develop Healthy Social Skills and Relationships: What the Research Shows About Navigating Adolescence*. Washington, DC: Child Trends; 2002
72. Resnick MD. Protective factors, resiliency and healthy youth development. *Adolesc Med*. 2000;11:157-165
73. Resnick MD. Resilience and protective factors in the lives of adolescents. *J Adolesc Health*. 2000;27:1-2
74. Steinberg L. Gallagher lecture. The family at adolescence: transition and transformation. *J Adolesc Health*. 2000;27:170-178
75. Steinberg L, Lamborn SD, Dornbusch SM, Darling N. Impact of parenting practices on adolescent achievement: authoritative parenting, school involvement, and encouragement to succeed. *Child Dev*. 1992;63:1266-1281

